

Literatur

Betrán AP, Meriáldi M, Lauer JA, Bing-Shun W, Thomas J, Van Look P, Wagner M: Rates of caesarean section: analysis of global, regional and national estimates. In: Paediatric and Perinatal Epidemiology 2007; 21: 98–113

Catling-Paull C, Johnston R, Ryan C, Foureur M, Homer C (2011): Non-clinical interventions that increase the uptake and success of vaginal birth after caesarean section: a systematic review. In: Journal of advanced Nursing 67(8), 1662–1676

Caughey A, Sundaram V, Kaimal AJ, Gienger A, Cheng Y, McDonald K, Shaffer B, Owens D, Bravata D: Systematic Review: Elective Induction of Labor Versus Expectant Management of Pregnancy. In: Ann Intern Med. 2009; 151:252-263

Chaillet N, Dumont A (2007): Evidence-Based Strategies for Reducing Cesarean Section Rates: A Meta-Analysis. In: Birth 34:1; 53

Hodnett ED, Gates S, Hofmeyr GJ, Sakala C, Weston J. Continuous support for women during childbirth. Cochrane Database of Systematic Reviews 2011, Issue 2. Art. No.: CD003766

Huh S Y, Rifas-Shiman S L, Zera C A, Rich Edwards J W, Oken E., Weiss S T, Gillman M W: Delivery by caesarean section and risk of obesity in pre-school age: a prospective cohort study. In: British Medical Journal 2012

Lumbiganon, Pisake, Laopaiboon, Malinee, Gülmezoglu, metin, Souza, Joao Paulo et al.: Method of delivery and pregnancy outcomes in Asia: the WHO global survey on maternal and perinatal health 2007-08, The Lancet 2010; 375(9713): 490-99

Lutz, Ulrike, Kolip, Petra, Die GEK-Kaiserschnitt-Studie (2006), http://www.ipp.uni-bremen.de/downloads/abteilung2/projekte/GEK_Kaiserschnittstudie.pdf

NICE (National Institute for Health and clinical Excellence) (2012): Women could avoid having unnecessary caesareans, [http://www.nice.org.uk/news/Women Could Avoid Having Unnecessary Caesarean](http://www.nice.org.uk/news/Women_Could_Avoid_Having_Unnecessary_Caesarean) (aufgerufen 3.4.2012)

Niino, Y: The increasing caesarean rate globally and what we can do about it. In: BioScience Trends 2011; 5(4): 139-150

Scarella A, Chamy V, Sepúlveda M, Belizán J (2011): Medical audit using the Ten Group Classification System and its impact on the caesarean section rate. In: European Journal of Obstetrics & Gynecology and Reproductive Biology 154 (2010) 136–140

Souza JP et al.: Caesarean section without medical indications is associated with an increased risk of adverse short-term maternal outcomes: the 2004–2008 WHO Global Survey on Maternal and Perinatal Health. In: BMC Medicine, 2010; 8 (1):71

Stjernholm Y, Petersson K, Eneroth E: Changed indications for caesarean sections. In: Acta Obstetrica et Gynecologica. 2010; 89: 49–53

Van Dillen J, Lim F, Van Rijssel E: Introducing caesarean section audit in a regional teaching hospital in The Netherlands. In: European Journal of Obstetrics & Gynecology and Reproductive Biology 139. 2008; 151–156
Villar J, Carroli G, Zavaleta N, Donner A, Wojdyla D, Faundes A, Velazko A, Bataglia V, Langer A, Narváez A, Valladares E, Shah A, Campodónico L, Romero M, Reynoso S, Simônia de Padua K, Giordano D, Kublickas M, Acosta A, Survey on Maternal and Perinatal Health Research Group: Maternal and neonatal individual risks and benefits associated with caesarean delivery: multicentre prospective study. In: BMJ 2007; 335; 1025

Wagner M: Choosing caesarean section. In: Lancet 2000; 356: 1677–80
Wagner M: Fish can't see water: the need to humanize birth. In: International Journal of Gynecology & Obstetrics 2001; 75: 25-37
WHO (2000): Erklärung von München – Pflegendende und Hebammen – ein Plus für Gesundheit. http://www.euro.who.int/__data/assets/pdf_file/0008/53855/E93016G.pdf am 16.5.2012 [Abruf 16.5.2012]

WHO (2002): Strategic Directions for Strengthening Nursing and Midwifery Services. <http://whqlibdoc.who.int/publications/2002/924156217X.pdf> [Abruf 16.5.2012]

WHO (2009): Making pregnancy safer – Assessment tool for the quality of hospital care for mothers and newborn babies. 47- 50. http://www.euro.who.int/__data/assets/pdf_file/0008/98792/E93128.pdf [Abruf 16.5.2012]

WHO Policy brief (2010): Caesarean section without medical indication increases risk of short-term adverse outcomes for mothers. http://whqlibdoc.who.int/hq/2010/WHO_RHR_HRP_10.20_eng.pdf [Abruf 16.5.2012]