

## Literatur zum Artikel "Wie Kaffeekränzchen ohne Torte vor Demenz schützen"

- Chittur SV, Sangster-Guity N, McCormick PJ (2008): Histone deacetylase inhibitors: a new mode for inhibition of cholesterol metabolism. *BMC Genomics*. 29; 9:507.
- Cooney RV, Franke AA, Harwood PJ, Hatch-Pigott V, Custer LJ, Mordan LJ (1993): Gamma-tocopherol detoxification of nitrogen dioxide: superiority to alpha-tocopherol. *Proc Natl Acad Sci USA*; 90(5):1771-5.
- Costa J1, Lunet N, Santos C, Santos J, Vaz-Carneiro A (2010): Caffeine exposure and the risk of Parkinson's disease: a systematic review and meta-analysis of observational studies. *J Alzheimers Dis*; 20 Suppl 1:S221-38.
- Dai Q, Borenstein AR, Wu Y, Jackson JC, Larson EB (2006): Fruit and vegetable juices and Alzheimer's disease: the Kame project. *Am J Med*; 119(9): 751-759.
- DAIG (Deutsche Alzheimer Gesellschaft) (2012): Selbsthilfe Demenz. Das Wichtigste 1: Die Epidemiologie der Demenz. URL: [http://www.deutsche-alzheimer.de/fileadmin/alz/pdf/factsheets/FactSheet01\\_2012.pdf](http://www.deutsche-alzheimer.de/fileadmin/alz/pdf/factsheets/FactSheet01_2012.pdf) (25.06.2013).
- de la Monte SM (2012): Contributions of brain insulin resistance and deficiency in amyloid-related neurodegeneration in Alzheimer's disease. *Drugs*; 72(1): 49-66.
- Eskelinen M (2004): The Effects of Midlife Diet on Late-Life Cognition: An Epidemiological Approach. *Publications of the University of Eastern Finland. Dissertations in Health Sciences* 220. 2014. 123 p.
- Eskelinen MH, Ngandu T, Tuomilehto J, Soininen H, Kivipelto M (2009): Midlife coffee and tea drinking and the risk of late-life dementia: a population-based CAIDE study. *J Alzheimers Dis*; 16(1):85-91.
- Fiocco AJ, Shatenstein B, Ferland G, Payette H, Belleville S, Kerfoot MJ, Morais JA, Greenwood CE (2012): Sodium intake and physical activity impact cognitive maintenance in older adults: the NuAge Study. *Neurobiol Aging*; 33(4): 829-e21-e28.
- Giem P, Beeson WL, Fraser GE (1993): The incidence of dementia and intake of animal products: preliminary findings from the adventist health study. *Neuroepidemiology*. 12:28-36.
- Gu Y, Nieves JW, Stern Y, Luchsinger JA, Scarmeas N (2010): Food combination and Alzheimer disease risk: a protective diet. *Arch Neurol*; 67(6): 699-706.
- Gustaw-Rothenberg K (2009): Dietary patterns associated with Alzheimer's disease: population based study. *Int J Environ Res Public Health*; 6(4): 1335-1340.
- Hartman RE, Shah A, Fagan AM, Schwetye KE, Parsadanian M, Schulman RN, Finn MB, Holtzman DM (2006): Pomegranate juice decreases amyloid load and improves behavior in a mouse model of Alzheimer's disease. *Neurobiol Dis*; 24(3): 506-515.
- Jiang Q, Christen S, Shigenaga MK, Ames BN (2001): gamma-tocopherol, the major form of vitamin E in the US diet, deserves more attention. *Am J Clin Nutr*; 74(6):714-22.
- Jiang Q, Elson-Schwab I, Courtemanche C, Ames BN (2000): gamma-tocopherol and its major metabolite, in contrast to alpha-tocopherol, inhibit cyclooxygenase activity in macrophages and epithelial cells. *Proc Natl Acad Sci USA*; 97(21):11494-9.
- Jomova K, Vondrakova D, Lawson M, Valko M (2010): Metals, oxidative stress and neurodegenerative disorders. *Mol Cell Biochem*; 345(1-2): 91-104.
- Kamal-Eldin A, Appelqvist LA (1996): The chemistry and antioxidant properties of tocopherols and tocotrienols. *Lipids*; 31(7):671-701.
- Kontush K, Schekatolina S (2004): Vitamin E in neurodegenerative disorders: Alzheimer's disease. *Ann NY Acad Sci*; 1031:249-62.
- Li FJ, Shen L, Ji HF (2012): Dietary intakes of vitamin E, vitamin C, and β-carotene and risk of Alzheimer's disease: a meta-analysis. *J Alzheimers Dis*; 31(2): 253-258.
- Luchsinger JA, Tang M, Shea S, Mayeux R (2002): Caloric intake and the risk of Alzheimer disease. *Arch Neurol*. 59:1258-1263.
- Martins IJ, Binosha Fernando WMAD (2014): High Fibre Diets and Alzheimer's Disease. *Food and Nutrition Sciences*, 5, 410-424
- Martins IJ, Hone E, Foster JK, Sünram-Lea SI, Gnjeć A, Fuller SJ, Nolan D, Gandy SE, Martins RN (2006): Apolipoprotein E, cholesterol metabolism, diabetes, and the convergence of risk factors for Alzheimer's disease and cardiovascular disease. *Mol Psychiatry*; 11(8): 721-736.
- Mertin A (2013): So schraubt Fett am ganzen Körper. Spiegel Online. URL: <http://www.spiegel.de/gesundheit/ernaehrung/wie-sich-fett-auf-den-stoffwechsel-und-die-gene-auswirkt-a-897441.html> (29.05.2013).
- Morris MC, Evans DA, Bienias JL, Tangney CC, Bennett DA, Aggarwal N, Schneider J, Wilson RS (2003): Dietary fats and the risk of incident Alzheimer disease. *Arch Neurol*; 60:194-200.
- Morris MC, Evans DA, Tangney CC, Bienias JL, Wilson RS, Aggarwal NT, Scherr PA (2005): Relation of the tocopherol forms to incident Alzheimer disease and to cognitive change. *Am J Clin Nutr*; 81(2):508-14.

- Morris MC, Schneider JA, Li H, Tangney CC, Nag S, Bennett DA, Honer WG, Barnes LL (2014): Brain tocopherols related to Alzheimer's disease neuropathology in humans. *Alzheimers Dement*; doi: 10.1016/j.jalz.2013.12.015. [Epub ahead of print].
- Mutter J, Naumann J, Schneider R, Walach H (2007): Quecksilber und Alzheimer-Erkrankung. *Fortschr Neurol Psychiatr*; 75(9): 528-538.
- Okereke OI, Rosner BA, Kim DH, Kang JH, Cook NR, Manson JE, Buring JE, Willett WC, Grodstein F (2012): Dietary fat types and 4-year cognitive change in community-dwelling older women. *Ann Neurol*; 72(1): 124-134.
- Ortega RM, Requejo AM, Andres P, López-Sobaler AM, Quintas ME, Redondo MR, Navia B, Rivas T (1997): Dietary intake and cognitive function in a group of elderly people. *Am J Clin Nutr*; 66: 803-809.
- Ozawa M, Ninomiya T, Ohara T, Hirakawa Y, Doi Y, Hata J, Uchida K, Shirota T, Kitazono T, Kiyohara Y (2012): Self-reported dietary intake of potassium, calcium, and magnesium and risk of dementia in the Japanese: the Hisayama Study. *J Am Geriatr Soc*. 60(8): 1515-20.
- Rota C, Rimbach G, Minihane AM, Stoecklin E, Barella L (2005): Dietary vitamin E modulates differential gene expression in the rat hippocampus: potential implications for its neuroprotective properties. *Nutr Neurosci*; 8(1):21-9.
- Rothgang H, Iwansky S, Müller R, Sauer S, Unger R (2010): BARMER GEK Pflegereport 2010. Schwerpunktthema: Demenz und Pflege. In: BARMER GEK (Hrsg.): Schriftenreihe zur Gesundheitsanalyse, Band 5. Asgard-Verlag, St. Augustin.
- Smith SR (2009): A look at the low-carbohydrate diet. *N Engl J Med*; 361(23): 2286-2288.
- Tucker JM, Townsend DM (2005): Alpha-tocopherol: roles in prevention and therapy of human disease. *Biomed Pharmacother*; 59(7):380-7.
- USDA (United States Department of Agriculture) (2014): National Nutrient Database for Standard Reference, Release 26, The National Agricultural Library. URL: <http://ndb.nal.usda.gov/ndb/foods> (25.06.2014).
- Williamson KS, Gabbita SP, Mou S, West M, Pye QN, Markesberry WR, Cooney RV, Grammas P, Reimann-Philipp U, Floyd RA, Hensley K (2002): The nitration product 5-nitro-gamma-tocopherol is increased in the Alzheimer brain. *Nitric Oxide*; 6(2):221-7.
- Zhuo JM, Portugal GS, Kruger WD, Wang H, Gould TJ, Pratico D (2010): Diet-induced hyperhomocysteinemia increases amyloid-beta formation and deposition in a mouse model of Alzheimer's disease. *Curr Alzheimer Res*; 7(2): 140-149.